

An Update on the X-ray transient Narrow-Line Seyfert 1 galaxy WPVS 007: *Swift* observations of UV variability and persistence of X-ray faintness

Dirk Grupe¹

grupe@astro.psu.edu

, Patricia Schady^{1,2}, Karen M. Leighly³, Stefanie Komossa⁴, Paul T. O'Brien⁵, John A. Nousek¹

ABSTRACT

We report on the detection of UV variability and the persistence of X-ray faintness of the X-ray transient Narrow-Line Seyfert 1 galaxy WPVS 007 based on the first year of monitoring this AGN with *Swift* between 2005 October and 2007 January. WPVS 007 has been an unusual source. While being X-ray bright during the ROSAT All-Sky Survey it has been extremely faint in all following X-ray observations. *Swift* also finds this NLS1 to be X-ray faint and not detected in the *Swift* X-Ray Telescope at an 3σ upper limit of $2.6 \times 10^{-17} \text{ W m}^{-2}$ in the 0.3-10.0 keV band and confirms that the AGN is still in a low state. During the 2006 July and December observations with *Swift*'s UV-Optical Telescope (UVOT) the AGN became fainter by about 0.2 mag in the UV filters and by about 0.1 mag in V, B, and U compared with the 2005 October to 2006 January and 2006 September/October observations followed by a rebrightening in the 2007 January observation. This variability can be caused either by a change in the absorption column density and therefore the reddening in the UV, or by flux variations of the central engine. We also noticed that the flux in the UVOT filters agree with earlier measurements by the International Ultraviolet Explorer taken between 1993-1995, but spectra taken by the Hubble Space Telescope Faint Object Spectrograph show that WPVS 007 was fainter in the UV by a factor of at least 2 in 1996. The flat optical/UV spectrum suggests that some UV extinction is present in the spectrum, but that alone cannot at all account for the dramatic fading in the X-ray flux. Most likely we see a partial covering absorber in X-rays. Alternatively, the current X-ray emission seen from WPVS 007 may also be the emission from the host galaxy.

Subject headings: galaxies: active, galaxies: individual (WPVS 007)

¹Department of Astronomy and Astrophysics, Pennsylvania State University, 525 Davey Lab, University Park, PA 16802

²Mullard Space Science Laboratory, Holmbury St. Mary, Dorking, Surrey RH5 6NT, U.K.; email: ps@mssl.ucl.ac.uk

³Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, 440 West Brooks Street, Norman, OK 73019; email: leighly@nhn.ou.edu

⁴Max-Planck-Institut für extraterrestrische Physik, Giessenbachstr., D-85748 Garching, Germany; email: skomossa@mpe.mpg.de

⁵Department of Physics & Astronomy, University of Le-

1. Introduction

When observed at optical or lower energies, radio-quiet AGN appear to be rather stable and not highly variable. However, this picture changes dramatically when AGN are observed in X-rays. Flux variability by factors 2-3 on timescales of days to months is quite common among low- and high-luminosity AGN. An increasing number that vary by factors 10-30 has emerged in the last decade, including in intermediate-type Seyferts

icester, Leicester, LE1 7R, UK, email: pto@star.le.ac.uk

and Narrow-Line Seyfert 1 galaxies (NLS1s), and there is good evidence that a substantial part of that variability is caused by (cold) absorption, in terms of complete or partial covering. Interestingly, the highest amplitudes of variability have been detected from the cores of non-active galaxies in terms of transient flares, interpreted as tidal disruptions of stars by the black holes at the centers of these galaxies (e.g. Komossa et al. 2004, and references therein).

NLS1s are objects with extreme properties: they show the steepest X-ray spectra, strongest optical FeII emission and weakest [OIII] emission (e.g. Boller et al. 1996; Boroson & Green 1992; Laor et al. 1997; Grupe et al. 2004a). The most common explanation for their extreme properties is that they have relatively small black hole mass and a high Eddington ratio L/L_{edd} (e.g. Boroson 2002; Grupe et al. 2004a; Sulentic et al. 2000). NLS1s are also known to be objects with very strong X-ray variability (e.g. Leighly 1999).

The NLS1 WPVS 007 (1RXS J003916.6–511701, RBS 0088; $\alpha_{2000} = 00^{\text{h}}39^{\text{m}}15.^{\text{s}}8$, $\delta_{2000} = -51^{\circ}17'03''0$, $z=0.029$; Grupe et al. 1995) was discovered in a survey of faint southern galaxies with H α emission by Wamsteker et al. (1985) and is a unique X-ray transient AGN. While X-ray transience is typically associated with an X-ray outburst caused by a dramatic increase in the accretion rate or the very onset of accretion, the situation is very different in WPVS 007. During the ROSAT All-Sky Survey (RASS, Voges et al. 1999), when the source was X-ray bright, the optical-to-X-ray flux ratio was in a normal range for AGN (e.g. Beuermann et al. 1999; Maccacaro et al. 1988). However, all follow-up X-ray observations between 1994 and 2002 using ROSAT and *Chandra* found it to have almost vanished from the X-ray sky (Grupe et al. 2001; Vaughan et al. 2004).

Until recently the cause for the transient behavior of WPVS 007 had been a mystery. Grupe et al. (1995) suggested that this transience could be due to a temperature change in the accretion disk that would shift the soft X-ray spectrum out of the ROSAT PSPC energy observing window. However, in recent years it became clear that the cause of the transience is absorption. In 1996 July, WPVS 007 was observed by HST (Goodrich 2000; Constantin & Shields 2003); a 2003 *FUSE* observation revealed the

emergence of a BAL flow (Leighly et al. 2005; Leighly et al., in preparation 2007). A discovery of a re-brightening and following rise in the X-ray luminosity, and in spectral changes, will set tight constraints on the movement and location of the absorber, and on the nature of the absorption. The high amplitude of the variability makes WPVS 007 exceptional among the known cases of absorption variability. In order to detect a possible re-brightening and therefore the disappearance of the absorber, we began a monitoring campaign using *Swift* in 2005 October.

The *Swift* mission (Gehrels et al. 2004) was launched on 2004 November 20th. The main purpose is to hunt and observe Gamma-Ray Bursts (GRBs). However, part of the observing time is used for fill-in targets and targets-of-opportunity when no GRB can be observed. Due to its multi-wavelength capacities and its flexible scheduling, *Swift* is an ideal observatory of all types of AGN, as demonstrated by e.g. Grupe et al. (2006) on the NLS1 RX J0148.3–2758 and Markwardt et al. (2005) on the search for obscured AGN in the BAT survey. *Swift* is equipped with three telescopes: at the high energy end the Burst Alert Telescope (BAT, Barthelmy 2005) operating in the 15-150 keV energy range, the X-Ray Telescope (XRT, Burrows et al. 2005), which covers the soft X-ray range between 0.3-10.0 keV, and at the long wavelength end, the UV-Optical Telescope (UVOT, Roming et al. 2005). The XRT uses a CCD detector identical to the EPIC MOS on-board XMM (Turner et al. 2001). The UVOT covers the range between 1700-6500Å and is a sister instrument of XMM's Optical Monitor (OM, Mason et al. 2001). The UVOT has a similar set of filters as the OM (Mason et al. 2001; Roming et al. 2005). However, the UVOT UV throughput is a factor of about 10 higher than in the OM.

The outline of this paper is as follows: in §2 we describe the *Swift* observations and the data reduction, in §3 we present the results of the *Swift* XRT and UVOT data analysis and compare the UVOT data with earlier IUE and HST spectra, and in §4 we discuss the results. Throughout the paper spectral indexes are denoted as energy spectral indexes with $F_{\nu} \propto \nu^{-\alpha}$. Luminosities are calculated assuming a Λ CDM cosmology with $\Omega_{\text{M}}=0.27$, $\Omega_{\Lambda}=0.73$ and a Hubble constant

of $H_0=75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ using a luminosity distances $D=118 \text{ Mpc}$ given by Hogg (1999). All errors are 1σ unless stated otherwise.

2. Observations and data reduction

WPVS 007 has been monitored by *Swift* between 2005 October and 2007 January. Table 1 lists the *Swift* XRT observations, including the start and end times, the total exposure times, and the 3σ upper limits. Note that we do not include the segment 005 data (2006 September 06) in the XRT analysis because during the time of that observation the XRT detector was rather warm resulting in an enhanced detector background. However, the UVOT data from that time period were not affected. The *Swift* UVOT observations are summarized in Table 2. Also note that segment 008 does not exist. WPVS 007 was originally scheduled for 2006 December 03, but the observations were superseded by the detections of GRBs 061201 and 061202 (Marshall et al. 2006; Sakamoto et al. 2006, respectively) before the start of the WPVS 007 observations. Segment numbers, however, can only be used once by the *Swift* scheduling tool. The segment numbers in both table refer to the days *Swift* observed WPVS 007 (See the description in Grupe et al. 2006). In the first observation of 2006 December (segment 009) we noticed that the AGN became significantly fainter in the UV by 0.2 mag. In order to investigate this behavior and to get a better estimate of the time scale we initiated an additional ToO for two pointings of 2 ks each which were executed on 2006 December 12 and 21 (segments 010 and 011). Also note that the 2007 January observation was split into two segments due to scheduling constraints, even though the observations were performed in consecutive orbits.

The XRT was operating in photon counting mode (Hill et al. 2004) and the data were reduced by the task *xrtpipeline* version 0.10.4., which is included in the HEASOFT package 6.1. Photons were collected in the 0.3-10.0 keV energy range. The upper limits were determined from the background in the XRT. The photons were extracted with *XSELECT* version 2.4. In order to compare the observations from different missions we use the HEASARC tool *PIMMS* version 3.8. For the conversion we assumed an absorbed power law model

with the absorption column density at the Galactic value ($2.84 \times 10^{20} \text{ cm}^{-2}$, Dickey & Lockman 1990) and an energy spectra slope $\alpha_X=3.0$. Note, however, that this is just an estimate, since we do not know what the low-state X-ray spectrum of WPVS 007 really looks like. For all observations the counts were corrected due to the exposure maps.

As listed in Table 2 UVOT observations were performed in all filters except during the 2005 October and December, and 2006 January observations only the UV filters were used. Photometry on all UVOT individual and coadded exposures was performed with the tool *uvotmaghist* version 0.1. A $6''$ and $12''$ radius extraction regions were used centered on WPVS 007 for the optical (V, B, and U) and UV filters (UV W1, UV M2, and UV W2), respectively and the background count rate was measured with a $20''$ radius aperture in a nearby source free region. All reported magnitudes have been corrected for Galactic extinction, where the reddening in the line-of-sight to the object is $E(B-V) = 0.012 \text{ mag}$ (Schlegel et al. 1998). All data were aspect corrected and coadded before measuring the magnitudes and fluxes. The UVOT uses Vega-based magnitudes with the following zeropoints: $V=17.88\pm0.09$, $B=19.16\pm0.12$, $U=18.38\pm0.23$, $UVW1=17.69\pm0.20$, $UVM2=17.29\pm0.23$, and $UVW2=17.77\pm0.20$ (Brown et al. 2007).

Prior the *Swift* observations WPVS 007 was observed in the UV by IUE four times between 1993 and 1995, HST in 1996 July, and FUSE in 2003 November. In this paper we make use of the IUE and HST data. Table 3 lists these observations. The FUSE data will be presented by Leighly et al., in preparation (2007) which will also contain a spectral analysis of the miniBALs present in the HST data.

3. Results

3.1. X-rays

We do not detect WPVS 007 in X-rays in any of the monitoring observations performed by *Swift* so far as listed in Table 1. To determine 3σ upper limits we applied the method by Kraft et al. (1991). This method determines the confidence levels for low numbers of counts using the Bayesian method for Poisson-distributed data. The back-

ground in all three observations was measured in a circular region with $r=235''$ around the position of WPVS 007. For the source itself we assumed an extraction radius of $23.5''$. The 3σ upper limits are listed in Table 1. We coadded all XRT observations together, except for the 2006 September 06 observation (segment 005) when the background was too high. From these coadded data with a total exposure time of 23.2 ks we measured an upper limit of 1.04×10^{-3} counts s^{-1} in the *Swift*-XRT. Assuming a power law model spectrum with $\alpha_X=3.0$ and N_H at the Galactic value this upper limit count rate converts to an upper limit in unabsorbed flux in the 0.3-10 keV band of 2.6×10^{-17} W m^{-2} .

Figure 1 displays the long term light curve of WPVS 007. The ROSAT values were taken from Grupe et al. (2001) and the *Chandra* data point from Vaughan et al. (2004). The light curve shows that WPVS 007 is still in a low state and the upper limit of the coadded *Swift* observations is consistent with the ROSAT PSPC and HRI detections. Note that Figure 1 uses PSPC counts which were converted by PIMMS assuming a power law spectrum with $\alpha_X=3.0$ and N_H at the Galactic value.

3.2. UVOT Photometry

The magnitudes in the UVOT filters are given in Table 2 and previous UV observations by IUE and HST are listed in Table 3. The errors quoted in Table 2 are statistical errors. As listed in Table 2 and shown in Figure 2 WPVS 007 became fainter in the optical filters by about 0.1 mag and by about 0.2 mag in the UV filters in the 2006 July and December observations compared with the observations obtained in 2005 and 2006 January and 2006 September and October. The AGN became brighter again by about 0.2 mag in the 2007 January observation which was about three weeks after the last observation in 2006 December. From the current light curve as shown in Figure 2 the AGN seems to be variable in the optical/UV on timescales of a few weeks to several months.

In order to determine whether this variability is real or just within the uncertainties we picked 4 field stars of similar magnitude as WPVS 007 as reference stars and compared their magnitudes segment by segment. Table 4 lists these 4 stars with the coordinates and their magnitudes in U,

UV W1, UV M2, and UV W2 with statistical errors. Figure 3 displays where these stars are located relative to WPVS 007. Note that during some of the observations not all the 4 stars were in the field of view of the UVOT. U magnitudes could only be given for Stars #3 and 4, because #1 and 2 are too bright in the U filter that they suffer significantly from coincident losses. Also note that star #1 and #2 were not in the field of view during some of the observations. As shown in Table 4, the variance in the field stars is small compared with the variability observed in WPVS 007. Therefore, we consider the UV variability found in WPVS 007 to be real. The change by 0.2 mag in the UV filters observed during the 2006 July and December observations is larger than the uncertainties between the measurements in the field stars.

WPVS 007 has shown variability on timescales of months before between the IUE and HST observations between 1993 and 1996. Figure 4 displays the *Swift* UVOT measurements from 2006 January, the HST spectrum from 1996 (Goodrich 2000; Constantin & Shields 2003), and the IUE spectrum averaging three spectra taken between 1993 and 1995. We excluded the 1995 November observation due to strong cosmic ray events during that observation. The data shown in Figure 4 are the observed data, uncorrected for reddening. The IUE spectra and the *Swift*-UVOT data seem to agree. However, the HST spectra show a significantly lower flux than the IUE and *Swift* data. Dunn et al. (2006) presented an Internet database¹ of UV continuum light curves of Seyfert galaxies which also showed that WPVS 007 displays significant variability in the UV between the IUE and HST observations. One possibility of a lower flux in the HST data is a mis-alignment of the source in the $1''$ aperture in the HST FOS. However, as listed in Table 3, the AGN was observed during several orbits and the fluxes in all these spectra agree with each other. A $1''$ aperture is also rather large compared to the $0''.1$ resolution. Therefore we exclude a mis-alignment as the cause of the lower flux in WPVS 007 during the HST observation in 1996 July. Note that Winkler et al. (1992) reports $V=15.28 \pm 0.03$, $B=15.77 \pm 0.03$, and $U=15.15 \pm 0.03$. While the small differences in V and B can be explained by

¹<http://www.chara.gsu.edu/PEGA/IUE>

the different central wavelength between the filters used by Winkler et al. (1992) and the UVOT, the difference in U suggests that WPVS 007 is variable in U.

3.3. Spectral Energy Distribution

Figure 5 displays the spectral energy distribution of WPVS 007 using 2MASS NIR data derived from the Nasa Extragalactic database, optical/UV data from *Swift*'s UVOT from 2006 July and the X-ray data from the *Chandra* observation from 2002 (Vaughan et al. 2004). The *Chandra* data are displayed as an un-absorbed power law model with $\alpha_X=3.0$. The optical-to-X-ray slope α_{ox}^2 measured from this plot is $\alpha_{ox}=5.4$. This is an extreme value for an AGN which typically have values around $\alpha_{ox}=1.5$ (e.g. Yuan et al. 1998; Strateva et al. 2005). Assuming an $\alpha_{ox}=1.5$ we would expect a flux at 2 keV $F_{2keV} = 6 \times 10^{-16} \text{ W m}^{-2}$, or a luminosity at 2 keV $L_{2keV} = 1 \times 10^{35} \text{ W}$. However, as shown in Figure 5, this is not the case. During the RASS observation, assuming an UV spectrum like during the *Swift* observations, the α_{ox} was in the order of $\alpha_{ox}=5.0$. The flattening of the UV spectrum, as shown by the UVOT photometry data, suggests some intrinsic reddening of the AGN. A NLS1 typically has a very blue optical/UV spectrum, as shown e.g. in Grupe et al. (2006) for the NLS1 RX J0148.3-2758. Assuming that the intrinsic optical UV spectrum of WPVS 007 is similar to that of RX J0148.3-2758, we can estimate a reddening by 0.6 mag in the UV W2 filter. This results in a lower limit of the intrinsic reddening $E_{B-V}=0.073$. Note that this is a rough estimate. Based on the $H\alpha/H\beta$ flux ratio, Winkler et al. (1992) estimated the extinction to $A_V \approx 1.0$ which is significantly higher than our estimate.

4. Discussion

Our main results are that WPVS 007 is still in a low state in X-rays and that it shows significant variability in the UV on timescales of months. Adding all *Swift* observations together (except for the segment 005 data of 2006 September 06) we determine a 3σ upper limit $= 1.04 \times 10^{-3}$ counts

s^{-1} , which converts to an unabsorbed flux in the 0.3-10 keV band of $2.6 \times 10^{-17} \text{ W m}^{-2}$. This upper limit is at a similar level to the ROSAT pointed PSPC and HRI observations. Note that *Chandra*, with its superb point spread function, was able to detect WPVS 007 at an even lower level (Vaughan et al. 2004) within an exposure time of 10 ks. Even though we could not detect a re-brightening of WPVS 007 in X-rays, it is still an exciting AGN. Note that the purpose of the *Swift* observations is not to obtain a deep detection, but to monitor the AGN in order to detect when it re-brightens again.

Our UVOT observations suggest that the AGN is variable in the UV bands by about 0.2 mag within timescales of a few months. UV variability, however, is not uncommon in AGN and has been reported for various AGN such as NGC 4151 (Edelson et al. 1996; Crenshaw, et al. 1996), NGC 5548 (Clavel et al. 1991; Korista et al. 1995), Akn 564 (Collier et al. 2001), and 3C 390.3 (O'Brien et al. 1998), but only a few NLS1s have repeated UV coverage as good as WPVS 007. In the case of WPVS 007 we can speculate that the UV variability is likely to be caused by a change in the absorption column density and therefore the reddening in the UV. Using the change in the UV W1 magnitude by 0.2 mag we can derive an additional $E_{B-V}=0.032$. This would cause an additional reddening by 0.10 mag in V, 0.17 in U, and 0.28 in UV W2 which is consistent with the changes we observe during the 2006 July and December observations. However, a change in luminosity of the central engine cannot completely be excluded based on the current data set.

The variability we observed in the UV between the IUE, HST and *Swift*-UVOT observations was previously also noticed by Dunn et al. (2006) between the IUE and HST observations. This is not a calibration issue. There are no problems reported on the HST FOS and IUE observations either. Also, looking at the light curves presented in the UV continuum light curve database by Dunn et al. (2006) suggest that the UV flux is already decaying between the IUE observation of 1994 to 1995 and that the 1996 HST observation is consistent with this decay.

WPVS 007 may be one of the most extreme cases of X-ray weak NLS1s such as those found by Williams et al. (2002, 2004). The reddened

²The X-ray loudness is defined by Tananbaum et al. (1979) as $\alpha_{ox} = -0.384 \log(f_{2keV}/f_{2500\text{\AA}})$.

optical/UV spectrum also suggests that the X-ray weakness of WPVS 007 is caused by absorption. However, of the 10 photons detected at the position of WPVS 007 in the *Chandra* ACIS-S data (Vaughan et al. 2004), 8 have energies below 1 keV. A simple cold absorber would have absorbed all these photons. The solution could be a partial covering absorber that would allow some of the soft X-ray photons to escape. Such partial covering absorbers have been found in several NLS1s, such as Mkn 1239 Grupe et al. (2004b) or 1H 0707-495 Gallo et al. (2004); Tanaka et al. (2004); Boller et al., (2002).

Alternatively, the low-state X-ray emission detected with *Chandra* may actually represent the X-ray emission from the host galaxy; the AGN being completely absorbed. In order to check this, we used the blue magnitude of the galaxy to predict the expected X-ray flux from the host galaxy, using the correlation between L_B and L_X for early-type galaxies of O’Sullivan et al., (2001). We use the extinction- corrected blue magnitudes measured with *Swift*, $m_{B,corr}=15.48$, and a previous USNO measurement, $m_{Bcorr}=14.24$.³ We then predict a host galaxy contribution to the X-ray luminosity of $L_X \approx 10^{33} - 7 \times 10^{33}$ W, which may well account for all of the observed low-state emission. Indeed, host galaxies typically show such soft, thermal spectra. However, as shown in Figure 1 in Vaughan et al. (2004) the 10 photons found at the position of WPVS 007 seem to be consistent with a point source. In order to verify this statement, a much longer observation by *Chandra* is needed. We will continue monitoring WPVS 007 every 4-6 weeks with *Swift* as long as the AGN is not sun-constrained. Especially more observations in the UV will give us a better handle on the timescales of the variability in the UV.

First we want to thank Neil Gehrels for approving our ToO requests and the *Swift* team for performing the ToO observations of WPVS 007 and scheduling the AGN on a regular basis. We would also like to thank Jay Dunn for quickly checking

the IUE and HST data for any problems, Matthias Dietrich for various discussions about UV variability in AGN, and the anonymous referee for useful comments and suggestions to improve the paper. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, Caltech, under contract with the National Aeronautics and Space Administration. This research was supported by NASA contract NAS5-00136 (D.G., & J.N.).

REFERENCES

- Barthelmy, S.D., 2005, *Space Science Reviews*, 120, 143
- Beuermann, K., Thomas, H.-C., Reinsch, K., et al., 1999, *A&A*, 347, 47
- Boller, T., Brandt, W.N., & Fink, H.H., 1996, *A&A*, 305, 53
- Boller, T., et al., 2002, *MNRAS*, 329, L1
- Boroson, T.A., & Green, R.F., 1992, *ApJS*, 80, 109
- Boroson, T.A., 2002, *ApJ*, 565, 78
- Brown, P.J., et al., 2007, *ApJ*, in press, astro-ph/0612541
- Burrows, D., et al., 2005, *Space Science Reviews*, 120, 165
- Clavel, J., et al., 1991, *ApJ*, 470, 322
- Collier, S., et al., 2001, *ApJ*, 561, 146
- Constantin, A., & Shields, J.C., 2003, *PASP*, 115, 592
- Crenshaw, D.M., et al., 1996, *ApJ*, 470, 322
- Dickey, J.M., & Lockman, F.J., 1990, *ARA&A*, 28, 215
- Dunn, J.P., Jackson, B., Deo, R.P., Farrington, C., Das, V., & Crenshaw, D.M., 2006, *PASP*, 118, 572
- Edelson, R.A., et al., 1996, *ApJ*, 470, 364
- Gallo, L.C., Tanaka, Y., Boller, T., Fabian, A.C., Vaughan, S., & Brandt, W.N., 2004, *MNRAS*, 353, 1064

³ Both measurements have to be taken with some caution, since we did not correct for any possible AGN contribution to m_B on the one hand, and since the short *Swift* observation may have missed part of the host galaxy contribution on the other hand. Therefore, the estimate should only be regarded as order of magnitude

- Gehrels, N., et al., 2004, *ApJ*, 611, 1005
- Goodrich, R.W., 2000, *New Astronomy reviews*, Vol 44, 519
- Grupe, D., 2004, *AJ*, 127, 1799
- Grupe, D., Beuermann, K., Mannheim, K., Thomas, H.-C., de Martino, D., & Fink, H.H., 1995, *A&A*, 300, L21
- Grupe, D., Wills, B.J., Wills, D., Beuermann, K., 1998, *A&A*, 333, 827
- Grupe, D., Thomas, H.-C., & Beuermann, K., 2001, *A&A*, 367, 470
- Grupe, D., Wills, B.J., Leighly, K.M., & Meusinger, H., 2004a, *AJ*, 127, 156
- Grupe, D., Mathur, S., & Komossa, S., 2004b, *AJ*, 127, 3161
- Grupe, D., Leighly, K.M., Komossa, S., Schady, P., O'Brien, P.T., Burrows, D.N., & Nousek, J.A., 2006, *AJ*, 132, 1189
- Hill, J.E., et al., 2004, *SPIE*, 5165, 217
- Hogg, D., 1999, *astro-ph/9905116*
- Komossa, S., Halpern, J., Schartel, N., & Hasinger, G., 2004, *ApJ*, 603, L17
- Korista, K.T., et al., 1995, *ApJS*, 97, 285
- Kraft, R.P., Burrows, D.N., & Nousek, J.A., 1991, *ApJ*, 374, 344
- Laor, A., Fiore, F., Elvis, M., Wilkes, B.J., & McDowell, J.C., 1997, *ApJ*, 477, 93
- Leighly, K.M., 1999, *ApJS*, 125, 297
- Leighly, K.M., Casebeer, D.A., Hamann, F., & Grupe, D., 2005, *AAS*, 207, 1804
- Leighly, K.M., et al., 2007, *ApJ*, to be submitted,
- Maccacaro, T., Gioia, I.M., Wolter, A., Zamorani, G., & Stocke, J.T., 1988, *ApJ*, 326, 680
- Markwardt, C.B., Tueller, J., Skinner, G.K., Gehrels, N., Barthelmy, S.D., & Mushotzky, R.F., 2005, *ApJ*, 633, L77
- Marshall, F.E., et al., 2006, *GCN report* 18.1
- Mason, K.O., et al., 2001, *A&A*, 365, L36
- Mathur, S., 2000, *MNRAS*, 314, L17
- O'Brien, P.T., et al., 1998, *ApJ*, 509, 163
- O'Sullivan E., Forbes D.A., Ponman T.J., 2001, *MNRAS* 328, 461
- Roming, P.W.A., et al., 2005, *Space Science Reviews*, 120, 95
- Sakamoto, T., et al., 2006, *GCN report* 19.1
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Strateva, I.V., Brandt, W.N., Schneider, D.P., Vanden Berk, D.G., & Vignali, C., 2005, *AJ*, 130, 387
- Sulentic, J.W., Zwitter, T., Marziani, P., & Dultzin-Hacyan, D., 2000, *ApJ*, 536, L5
- Tanaka, Y., Boller, Th., Gallo, L.C., Keil, R., & Ueda, Y., 2004, *PASJ*, 56, L9
- Tananbaum, H., et al., 1979, *ApJ*, 234, L9
- Turner, M.J.L., Abbey, A., Arnaud, M., et al., 2001, *A&A*, 365, L27
- Vaughan, S., Edelson, R., & Warwick, R.S., 2004, *MNRAS*, 349, L1
- Voges, W., Aschenbach, B., Boller, T., et al., 1999, *A&A*, 349, 389
- Wamsteker, W., Prieto A., Vitorres, A., Schuster, H.E., Danks, A.C., Gonzalez R., & Rodriguez, G., 1985, *A&AS*, 62, 255
- Williams, R.J., Pogge, R.W., & Mathur, S., 2002, *AJ*, 124, 3042
- Williams, R.J., Pogge, R.W., & Mathur, S., 2004, *ApJ*, 610, 737
- Winkler, H., Stirpe, G.M., & Sekiguchi, K., 1992, *A&AS*, 94, 103
- Yuan, W., Siebert, J., Brinkmann, W., 1998, *A&A*, 334, 498

This 2-column preprint was prepared with the AAS L^AT_EX macros v5.2.




Fig. 1.— Long-term light curve of WPVS 007 containing the ROSAT All-Sky Survey and pointed PSPC and HRI observations, the *Chandra*, and the upper limits derived from the *Swift* XRT observations. The count rates were converted by assuming an absorbed power law model with $N_{\text{H}}=2.84 \times 10^{20} \text{ cm}^{-2}$ and $\alpha_{\text{X}}=3.0$. The large red downward arrow displays the upper limit of the coadded exposures of the *Swift*-XRT observations (23.2 ks).




Fig. 2.— *Swift* UVOT light curves of WPVS 007. The values are given in Table 2.




Fig. 3.— *Swift* UVOT W2 image of the 2006 January observation with the reference stars as listed in Table 4 and WPVS 007.




Fig. 4.— UV observations of WPVS 007. The red points are the *Swift* UVOT observations from 2006 January, the black spectrum is the HST observation from 1996, and the green spectrum the average of the IUE observation from 1993, 1994, and 1995 December.




Fig. 5.— Spectral energy distribution of WPVS 007 using the NIR data from 2MASS, optical/UV from the *Swift* UVOT observation from January 2006 corrected for Galactic reddening, and the X-ray data from the *Chandra* observation from 2002. The Chandra data are displayed as a power law model with $\alpha_X=3.0$. Note that all these observations have not been performed simultaneously. The HST spectra and the observed RASS data are displayed in light green/grey for comparison purposes.

TABLE 1
Swift XRT OBSERVATION LOG OF WPVS 007

Segment	T-start ¹	T-stop ¹	T _{exp} ²	3 σ ul ³
001	2005-10-20 03:21	2005-10-20 03:54	2025	5.78
002	2005-12-07 13:29	2005-12-07 13:58	1696	3.42
003	2006-01-05 00:48	2006-01-05 05:41	3466	1.67
004	2006-07-31 04:39	2006-07-31 06:30	1610	6.16
005	2006-09-06 08:19	2006-09-06 11:38	917	— ⁴
006	2006-09-11 05:40	2006-09-11 07:37	2335	5.02
007	2006-10-27 02:34	2006-10-27 23:33	3228	1.80
009	2006-12-06 00:02	2006-12-06 06:36	2964	2.70
010	2006-12-12 00:33	2006-12-12 01:01	1611	3.60
011	2006-12-21 06:19	2006-12-21 08:15	2273	4.36
012+013	2007-01-10 21:21	2007-01-11 00:53	1905	3.05
001-013 ⁴	2005-10-20 03:21	2007-01-11 05:53	23198 ⁴	1.04 ⁴

¹Start and End times are given in UT

²Observing time given in s

³3 σ upper limits in units of 10⁻³ XRT counts s⁻¹

⁴Due high background during the observation the segment 005 XRT data were not included in the analysis. Segment 008 does not exist (see text)

TABLE 2
Swift UVOT OBSERVATION OF WPVS 007

Segment	V		B		U		UV W1		UV M2		UV W2	
	T _{exp} ¹	Magcorr ²	T _{exp} ¹	Magcorr ²	T _{exp} ¹	Magcorr ²	T _{exp} ¹	Magcorr ²	T _{exp} ¹	Magcorr ²	T _{exp} ¹	Magcorr ²
001	...	...	...	...	...	...	656	14.28±0.01	686	14.50±0.01	686	14.53±0.01
002	...	...	...	...	...	...	550	14.31±0.01	588	14.48±0.01	588	14.55±0.01
003	...	...	...	...	...	...	1056	14.27±0.01	1171	14.46±0.01	1171	14.56±0.01
004	155	15.14±0.03	159	15.48±0.02	159	14.44±0.02	319	14.49±0.02	118	14.63±0.04	615	14.82±0.01
005	55	15.01±0.05	170	15.38±0.02	170	14.38±0.02	340	14.27±0.02	144	14.40±0.03	392	14.52±0.02
006	194	15.04±0.03	194	15.37±0.02	194	14.30±0.02	387	14.26±0.01	536	14.42±0.01	777	14.49±0.01
007	344	15.03±0.02	336	15.39±0.01	335	14.33±0.01	686	14.26±0.01	767	14.47±0.01	767	14.49±0.01
009	245	15.12±0.02	245	15.52±0.02	245	14.43±0.02	486	14.44±0.01	621	14.63±0.02	978	14.72±0.01
010	134	15.08±0.03	134	15.46±0.03	134	14.58±0.02	267	14.45±0.01	376	14.64±0.02	534	14.79±0.01
011	187	15.15±0.02	187	15.55±0.02	187	14.52±0.02	374	14.45±0.02	513	14.68±0.02	750	14.81±0.01
012+013	155	15.11±0.03	155	15.40±0.02	155	14.35±0.02	312	14.35±0.02	373	14.48±0.02	625	14.60±0.01

¹Observing time given in s

²Magnitude corrected for reddening with $E_{B-V}=0.012$ given by Schlegel et al. (1998). The errors given in this table are statistical errors

TABLE 3
PREVIOUS UV OBSERVATIONS OF WPVS 007

Mission	ObsID	grating	T-start ¹	T _{exp} ²
IUE	SWP 48542	—	1993-09-05 15:58	16800
	SWP 52369	—	1994-10-10 15:30	18900
	SWP 56215	—	1995-11-19 22:00	20000
	SWP 56318	—	1995-12-20 12:47	18000
HST ³	FOS Y3790102T	G130H	1996-07-30 13:30	1730
	FOS Y3790103T	G130H	1996-07-30 14:46	2110
	FOS Y3790104T	G160L	1996-07-30 15:30	240
	FOS Y3790105T	G190H	1996-07-30 16:29	1500
	FOS Y3790107T	G270H	1996-07-30 18:02	1280

¹Start and End times are given in UT

²Observing time given in s

³A complete listing of all HST FOS observations is given in Constantin & Shields (2003).

TABLE 4

LIST OF REFERENCE STARS USED TO DETERMINE THE ERROR IN THE PHOTOMETRY. THE MAGNITUDES ARE THE UNCORRECTED, DIRECTLY MEASURED VALUES. THE POSITIONS RELATIVE TO WPVS 007 ARE SHOWN IN FIGURE 3.

Object	α_{2000}	δ_{2000}	Filter	Segment 003	Segment 004	Segment 006	Segment 007	Segment 009	Segment 010	Segment 011	Seg. 012+013
Star 1	00 39 20.0	-51 10 53.2	UV W1	13.50 $\pm$ 0.01	...	...	13.50 $\pm$ 0.01	13.51 $\pm$ 0.01	13.53 $\pm$ 0.01	13.52 $\pm$ 0.01	13.50 $\pm$ 0.01
			UV M2	16.27 $\pm$ 0.02	...	...	16.26 $\pm$ 0.03	16.24 $\pm$ 0.03	16.30 $\pm$ 0.04	16.20 $\pm$ 0.03	16.22 $\pm$ 0.04
			UV W2	15.32 $\pm$ 0.02	...	...	15.26 $\pm$ 0.01	15.02 $\pm$ 0.01	15.10 $\pm$ 0.02	15.29 $\pm$ 0.01	15.32 $\pm$ 0.02
Star 2	00 39 54.2	-51 15 04.7	UV W1	14.38 $\pm$ 0.01	14.40 $\pm$ 0.02	14.40 $\pm$ 0.01	...	...	14.39 $\pm$ 0.02	14.44 $\pm$ 0.02	...
			UV M2	15.77 $\pm$ 0.02	15.78 $\pm$ 0.06	15.80 $\pm$ 0.03	...	...	15.79 $\pm$ 0.03	15.77 $\pm$ 0.03	...
			UV W2	15.93 $\pm$ 0.02	15.98 $\pm$ 0.02	15.95 $\pm$ 0.02	...	...	15.98 $\pm$ 0.02	15.99 $\pm$ 0.02	...
Star 3	00 39 01.8	-51 18 17.6	U	...	13.21 $\pm$ 0.02	13.27 $\pm$ 0.02	13.26 $\pm$ 0.01	13.21 $\pm$ 0.02	13.29 $\pm$ 0.02	13.25 $\pm$ 0.02	13.21 $\pm$ 0.02
			UV W1	14.30 $\pm$ 0.01	14.32 $\pm$ 0.02	14.37 $\pm$ 0.01	14.33 $\pm$ 0.01	14.30 $\pm$ 0.01	14.35 $\pm$ 0.02	14.33 $\pm$ 0.02	14.32 $\pm$ 0.02
			UV M2	15.25 $\pm$ 0.01	15.24 $\pm$ 0.04	15.35 $\pm$ 0.02	15.36 $\pm$ 0.02	15.29 $\pm$ 0.01	15.38 $\pm$ 0.03	15.31 $\pm$ 0.02	15.27 $\pm$ 0.02
Star 4	00 39 18.7	-51 14 14.0	UV W2	15.50 $\pm$ 0.01	15.55 $\pm$ 0.02	15.65 $\pm$ 0.02	15.54 $\pm$ 0.01	15.55 $\pm$ 0.01	15.64 $\pm$ 0.02	15.53 $\pm$ 0.02	15.54 $\pm$ 0.02
			U	...	13.89 $\pm$ 0.02	13.87 $\pm$ 0.02	13.88 $\pm$ 0.02	13.93 $\pm$ 0.02	13.92 $\pm$ 0.02	13.96 $\pm$ 0.02	13.91 $\pm$ 0.02
			UV W1	15.06 $\pm$ 0.01	15.04 $\pm$ 0.02	15.04 $\pm$ 0.02	15.04 $\pm$ 0.01	15.08 $\pm$ 0.02	15.05 $\pm$ 0.02	15.04 $\pm$ 0.02	15.09 $\pm$ 0.03
			UV M2	16.17 $\pm$ 0.02	16.07 $\pm$ 0.06	16.08 $\pm$ 0.03	16.12 $\pm$ 0.02	16.07 $\pm$ 0.03	16.14 $\pm$ 0.04	16.14 $\pm$ 0.03	16.07 $\pm$ 0.03
			UV W2	16.50 $\pm$ 0.02	16.41 $\pm$ 0.03	16.43 $\pm$ 0.02	16.46 $\pm$ 0.02	16.48 $\pm$ 0.02	16.43 $\pm$ 0.03	16.47 $\pm$ 0.03	16.55 $\pm$ 0.03